

AD-A077 137

WISCONSIN UNIV-MADISON MATHEMATICS RESEARCH CENTER
NONLINEAR SCHROEDINGER EVOLUTION EQUATIONS.(U)
SEP 79 H BREZIS , T GALLOUET

F/G 12/1

UNCLASSIFIED

MRC-TSR-1992

DAA629-75-C-0024
NL

| OF |
AD
A077137



END
DATE
FILMED
12-79
DDC

LEVEL


MRC Technical Summary Report #1992 ✓

NONLINEAR SCHRÖDINGER EVOLUTION
EQUATIONS

H. Brezis and T. Gallouet

AD A 077137

Mathematics Research Center
University of Wisconsin-Madison
610 Walnut Street
Madison, Wisconsin 53706



September 1979

Received July 30, 1979)

DDC FILE COPY

See 1473
in
back

Approved for public release
Distribution unlimited

Sponsored by
U. S. Army Research Office
P. O. Box 12211
Research Triangle Park
North Carolina 27709

UNIVERSITY OF WISCONSIN-MADISON
MATHEMATICS RESEARCH CENTER

NONLINEAR SCHRÖDINGER EVOLUTION EQUATIONS

H. Brezis¹ and T. Gallouet

Technical Summary Report #1992
September 1979

ABSTRACT

We consider the nonlinear Schrödinger equation

$$\begin{cases} i \frac{\partial u}{\partial t} - \Delta u + k|u|^2 u = 0 & \text{in } \Omega \times [0, \infty) \\ u(x, t) = 0 & \text{in } \partial\Omega \times [0, \infty) \\ u(x, 0) = u_0(x) & \text{in } \Omega \end{cases}$$

where Ω is a bounded domain or an exterior domain of \mathbb{R}^2 . Such an equation has been extensively studied when $\Omega = \mathbb{R}^2$, but the methods do not apply if $\Omega \neq \mathbb{R}^2$. We prove that there exists a unique global smooth solution if $k \geq 0$ or if $k < 0$ and $|k| \int |u_0|^2 < 4$. The proof relies on a new interpolation-embedding inequality:

$$\|u\|_{L^\infty} \leq C[1 + \sqrt{\log(1 + \|u\|_{H^2})}] \quad \text{for every } u \in H^2 \text{ with } \|u\|_{H^1} \leq 1.$$

AMS (MOS) Subject Classifications: 47H15, 46E35, 35K55, 35L60, 45E30

Key Words: Nonlinear Schrödinger equation, Global solutions, Sobolev embedding, Interpolation inequality

Work Unit Number 1 (Applied Analysis)

¹Dept. de Mathématiques, Université Paris VI, 4, pl. Jussieu, 75230 Paris Cedex 05, France.

SIGNIFICANCE AND EXPLANATION

The nonlinear Schrödinger equation occurs in the study of some problems in nonlinear optics (propagation of laser beams through the atmosphere or in a plasma). ~~It has been considered by many authors.~~ The main novelty of the present result is that it applies to the propagation of beams in channels (instead of the whole space). The proof relies on a new Sobolev-Orlicz embedding inequality which could be useful in other situations.

Accession For	
NTIS GMA&I	<input checked="checked" type="checkbox"/>
DDC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist.	Avail and/or special
A	

The responsibility for the wording and views expressed in this descriptive summary lies with MRC, and not with the authors of this report.

NONLINEAR SCHRÖDINGER EVOLUTION EQUATIONS

H. Brezis¹ and T. Gallouet

Let Ω be a domain in \mathbb{R}^2 with compact smooth boundary Γ (Ω could be for example a bounded domain or an exterior domain). Consider the equation

$$(1) \quad \begin{cases} i \frac{\partial u}{\partial t} - \Delta u + k|u|^2 u = 0 & \text{in } \Omega \times [0, \infty) \\ u(x, t) = 0 & \text{in } \Gamma \times [0, \infty) \\ u(x, 0) = u_0(x), \end{cases}$$

where $u(x, t)$ is a complex valued function and $k \in \mathbb{R}$ is a constant. Problem (1) which occurs in nonlinear optics when $\Omega = \mathbb{R}^2$ has been extensively studied in this case (see [1], [2], [3], [5], [8]), but we are not aware of any known result when $\Omega \neq \mathbb{R}^2$.

Our main result is the following:

Theorem 1. Let $u_0 \in H^2(\Omega) \cap H_0^1(\Omega)$. Assume that one of the following conditions holds

- (a) either $k \geq 0$,
- (b) or $k < 0$ and $|k| \int |u_0(x)|^2 dx < 4$.

Then there exists a unique solution of (1) such that

$$u \in C([0, \infty); H^2(\Omega)) \cap C^1([0, \infty); L^2(\Omega)).$$

The proof of Theorem 1 relies on several Lemmas. The first Lemma is of interest for its own sake; it is a new interpolation-embedding inequality.

¹Dept. de Mathématiques, Université Paris VI, 4, pl. Jussieu, 75230 Paris Cedex 05, France.

In what follows we denote by C various constants depending only on Ω .

Lemma 2. We have

$$(2) \quad \|u\|_{L^\infty} \leq C(1 + \sqrt{\log(1 + \|u\|_{H^2})})$$

for every $u \in H^2(\Omega)$ with $\|u\|_{H^1} \leq 1$.

Proof. It is well known that an H^2 function on Ω can be extended by an H^2 function on \mathbb{R}^2 . More precisely one can construct an extension operator P such that:

P is a bounded operator from $H^1(\Omega)$ into $H^1(\mathbb{R}^2)$

P is a bounded operator from $H^2(\Omega)$ into $H^2(\mathbb{R}^2)$

$Pu|_{\Omega} = u$ for every $u \in H^1(\Omega)$.

Let $u \in H^2(\Omega)$ with $\|u\|_{H^1} \leq 1$. Let $v = Pu$ and denote by \hat{v} the Fourier transform of v . We clearly have

$$(3) \quad \|(1 + |\xi|) \hat{v}\|_{L^2(\mathbb{R}^2)} \leq C$$

$$(4) \quad \|(1 + |\xi|^2) \hat{v}\|_{L^2(\mathbb{R}^2)} \leq C\|u\|_{H^2(\Omega)}$$

$$(5) \quad \|u\|_{L^\infty(\Omega)} \leq \|v\|_{L^\infty(\mathbb{R}^2)} \leq C\|\hat{v}\|_{L^1(\mathbb{R}^2)}.$$

For $R > 0$ we write

$$\begin{aligned} \|\hat{v}\|_{L^1} &= \int_{|\xi| < R} |\hat{v}(\xi)| d\xi + \int_{|\xi| \geq R} |\hat{v}(\xi)| d\xi = \int_{|\xi| < R} (1 + |\xi|) |\hat{v}(\xi)| \frac{1}{1 + |\xi|} d\xi \\ &\quad + \int_{|\xi| \geq R} (1 + |\xi|^2) |\hat{v}(\xi)| \frac{1}{1 + |\xi|^2} d\xi \\ &\leq C \left[\int_{|\xi| < R} \frac{1}{(1 + |\xi|)^2} d\xi \right]^{1/2} + C\|u\|_{H^2} \left[\int_{|\xi| \geq R} \frac{1}{(1 + |\xi|^2)^2} d\xi \right]^{1/2} \end{aligned}$$

by Cauchy-Schwarz, (3) and (4). A straightforward computation leads to

$$\|\hat{v}\|_{L^1} \leq C[\log(1+R)]^{1/2} + C\|u\|_{H^2}(1+R)^{-1}$$

by every $R \geq 0$. We obtain (2) by choosing $R = \|u\|_{H^2}$.

Lemma 3. We have

$$(6) \quad \| |u|^2 u \|_{H^2} \leq C \|u\|_{L^\infty}^2 \|u\|_{H^2} \quad \text{for every } u \in H^2(\Omega).$$

Proof of Lemma 3. Let D denote any first order differential operator. For $u \in H^2$ we have

$$|D^2(|u|^2 u)| \leq C(|u|^2 |D^2 u| + |u| |Du|^2),$$

and so

$$(7) \quad \| |u|^2 u \|_{H^2} \leq C \|u\|_{L^\infty}^2 \|u\|_{H^2} + C \|u\|_{L^\infty} \|u\|_{W^{1,4}}^2.$$

On the other hand an inequality of Gagliardo-Nirenberg (see [6]) implies that

$$(8) \quad \|u\|_{W^{1,4}} \leq C \|u\|_{L^\infty}^{1/2} \|u\|_{H^2}^{1/2}.$$

Combining (7) and (8) we obtain (6).

Finally we recall the following well known result essentially due to Segal [7]:

Lemma 4. Assume H is a Hilbert space and $A : D(A) \subset H \rightarrow H$ is an m -accretive linear operator. Assume F is a mapping from $D(A)$ into itself which is Lipschitz on every bounded set of $D(A)$. Then for every $u_0 \in D(A)$, there exists a unique solution u of the equation

$$\begin{cases} \frac{du}{dt} + Au = Fu \\ u(0) = u_0 \end{cases}$$

defined for $t \in [0, T_{\max})$ such that

$$u \in C^1([0, T_{\max}); H) \cap C([0, T_{\max}); D(A))$$

with the additional property that

$$\begin{cases} \text{either } T_{\max} = \infty \\ \text{or } T_{\max} < \infty \text{ and } \lim_{t \uparrow T_{\max}} \|u(t)\| + \|\Delta u(t)\| = \infty. \end{cases}$$

Proof of Theorem 1. We apply Lemma 4 in $H = L^2(\Omega)$ to $\Delta u = i\Delta u$, $D(A) = H^2(\Omega) \cap H_0^1(\Omega)$, $Fu = ik|u|^2u$. We shall show that $T_{\max} = \infty$ by proving that $\|u(t)\|_{H^2}$ remains bounded on every finite time interval.

First we multiply (1) by \bar{u} and consider the imaginary part. This leads to

$$(9) \quad \|u(t)\|_{L^2}^2 = \|u_0\|_{L^2}^2.$$

Next we multiply (1) by $\frac{\partial \bar{u}}{\partial t}$ and consider the real part. This leads to

$$(10) \quad \frac{1}{2} \int |\nabla u(x,t)|^2 dx + \frac{k}{4} \int |u(x,t)|^4 dx \equiv E_0$$

where

$$E_0 = \frac{1}{2} \int_{\Omega} |\nabla u_0(x)|^2 dx + \frac{k}{4} \int_{\Omega} |u_0(x)|^4 dx.$$

We claim that $\|u(t)\|_{H^1}$ remains bounded for $t > 0$. Indeed, this is clear when $k \geq 0$. While if $k < 0$ we have

$$(11) \quad \int |\nabla u(x,t)|^2 \leq \frac{|k|}{2} \int |u(x,t)|^4 dx + 2E_0.$$

On the other hand an inequality of Gagliardo and Nirenberg ([6]) shows that⁽¹⁾

⁽¹⁾ In order to obtain the constant $1/2$ one proceeds as follows. For $\varphi \in C_0^\infty(\mathbb{R}^2)$ we have $|\varphi(x_1, x_2)| \leq \frac{1}{2} \int_{-\infty}^{+\infty} |\varphi_{x_1}(t, x_2)| dt$, $|\varphi(x_1, x_2)| \leq \frac{1}{2} \int_{-\infty}^{+\infty} |\varphi_{x_2}(x_1, s)| ds$. Thus $\int_{\mathbb{R}^2} |\varphi|^2 dx \leq \frac{1}{4} \int_{\mathbb{R}^2} |\varphi_{x_1}| dx \int_{\mathbb{R}^2} |\varphi_{x_2}| dx$. Choosing $\varphi = |u|^2$ leads to

$$\int |u|^4 dx \leq \int |u|^2 dx \left(\int |u_{x_1}|^2 dx \right)^{1/2} \left(\int |u_{x_2}|^2 dx \right)^{1/2} \leq \frac{1}{2} \int |u|^2 dx \int |\nabla u|^2 dx.$$

$$\begin{aligned}
 (12) \quad \int |u|^4 dx &\leq \frac{1}{2} \int |u|^2 dx \int |\nabla u|^2 dx \\
 &= \frac{1}{2} \int |u_0|^2 dx \int |\nabla u|^2 dx .
 \end{aligned}$$

Combining (11), (12) and assumption (b) in Theorem 1 we see that

$$(13) \quad \|u(t)\|_{H^1} \leq C$$

where C is independent of t .

We now denote by $S(t)$ the L^2 isometry group generated by $-A$. From (1) we have

$$u(t) = S(t)u_0 + ik \int_0^t S(t-s)|u(s)|^2 u(s) ds$$

and so

$$Au(t) = S(t)Au_0 + ik \int_0^t S(t-s)A[|u(s)|^2 u(s)] ds .$$

Thus

$$(14) \quad \|Au(t)\|_{L^2} \leq \|Au_0\|_{L^2} + |k| \int_0^t \|A[|u(s)|^2 u(s)]\|_{L^2} ds .$$

Lemma 3 implies that

$$\|A[|u(s)|^2 u(s)]\|_{L^2} \leq C \|u(s)\|_{L^\infty}^2 \|u(s)\|_{H^2} .$$

From Lemma 2 and estimate (13) we deduce that

$$\|u(s)\|_{L^\infty} \leq C(1 + \sqrt{\log(1 + \|u(s)\|_{H^2})}) .$$

Hence (14) leads to

$$(15) \quad \|u(t)\|_{H^2} \leq C + C \int_0^t \|u(s)\|_{H^2} [1 + \log(1 + \|u(s)\|_{H^2})] ds .$$

We denote by $G(t)$ the RHS in (15); thus

$$G'(t) = C \|u(t)\|_{H^2}^2 [1 + \log(1 + \|u(t)\|_{H^2}^2)] \leq CG(t) [1 + \log(1 + G(t))] .$$

Consequently

$$\frac{d}{dt} \log[1 + \log(1 + G(t))] \leq C$$

and we find an estimate for $\|u(t)\|_{H^2}$ of the form

$$\|u(t)\|_{H^2} \leq e^{\alpha} e^{\beta t}$$

for some constants α and β . Therefore $\|u(t)\|_{H^2}$ remains bounded on every finite time interval and so we must have $T_{\max} = \infty$.

Remarks. 1) The proof of Theorem 1 leads to an estimate of the form

$\|u(t)\|_{L^\infty} \leq \alpha e^{\beta t}$. We do not know whether $\|u(t)\|_{L^\infty}$ remains actually bounded as $t \rightarrow \infty$.

2) When $k < 0$ and $|k| \int |u_0|^2 > 4$, it is known (see [4],[2]) if $\Omega = \mathbb{R}^2$ that the solution of (1) corresponding to some initial conditions may blow up in finite time. A similar phenomenon presumably occurs when $\Omega \neq \mathbb{R}^2$.

REFERENCES

- [1] J. B. Baillon, T. Cazenave, and M. Figueira, Equation de Schrödinger nonlinéaire, C. R. Acad. Sc. Paris 284 (1977), p. 869-872.
- [2] T. Cazenave, Equations de Schrödinger nonlinéaires,
- [3] J. Ginibre and G. Velo, On a class of nonlinear Schrödinger equations,
- [4] R. T. Glassey, On the blowing up of solutions to the Cauchy problem for the nonlinear Schrödinger equation, J. Math. Phys. 18 (1977), p. 1794-1979.
- [5] J. E. Lin and W. A. Strauss, Decay and scattering of solutions of a nonlinear Schrödinger equation, J. Funct. Anal. 30 (1978), p. 245-263.
- [6] L. Nirenberg, On elliptic partial differential equations, Ann. Sc. Norm. Sup. Pisa 13 (1959), p. 115-162.
- [7] I. Segal, Nonlinear semi-groups, Ann. of Math. 78 (1963), p. 339-364.
- [8] W. A. Strauss, The nonlinear Schrödinger equation, in Contemporary Developments in Continuum Mechanics and PDE, G. de la Penha and L. Medeiros ed., North Holland (1978), p. 452-465.

HB/TG/scr

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 1992	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) NONLINEAR SCHRÖDINGER EVOLUTION EQUATIONS		5. TYPE OF REPORT & PERIOD COVERED Summary Report - no specific reporting period
7. AUTHOR(s) H. Brezis and T. Gallouet		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Mathematics Research Center, University of Wisconsin 610 Walnut Street Madison, Wisconsin 53706		8. CONTRACT OR GRANT NUMBER(s) 15 DAAG29-75-C-0024
11. CONTROLLING OFFICE NAME AND ADDRESS U. S. Army Research Office P. O. Box 12211 Research Triangle Park, North Carolina 27709		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS Work Unit Number 1 - Applied Analysis
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) 9 Technical summary rept.		12. REPORT DATE 11 September 1979
		13. NUMBER OF PAGES 7
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited. 14 MRC-TSR-1992		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Nonlinear Schrödinger equation Global solutions Sobolev embedding Interpolation inequality		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) We consider the nonlinear Schrödinger equation $\begin{cases} i \frac{\partial u}{\partial t} - \Delta u + k u ^2 u = 0 & \text{in } \Omega \times [0, \infty) \\ u(x, t) = 0 & \text{in } \partial\Omega \times [0, \infty) \\ u(x, 0) = u_0(x) & \text{in } \Omega \end{cases}$		

20. ABSTRACT - Cont'd.

where Ω is a bounded domain or an exterior domain of \mathbb{R}^2 . Such an equation has been extensively studied when $\Omega = \mathbb{R}^2$, but the methods do not apply if $\Omega \neq \mathbb{R}^2$. We prove that there exists a unique global smooth solution if $k \geq 0$ or if $k < 0$ and $|k| \int |u_0|^2 < 4$. The proof relies on a new interpolation-embedding inequality:

$$\|u\|_{L^\infty} \leq C(1 + \sqrt{\log(1 + \|u\|_{H^2})}) \quad \text{for every } u \in H^2 \text{ with } \|u\|_{H^1} \leq 1.$$